

CONFIRMATION OF ERRORS IN *HIPPARCOS* PARALLAXES FROM HST/FGS ASTROMETRY OF THE PLEIADES*

DAVID R. SODERBLOM AND ED NELAN

Space Telescope Science Institute², 3700 San Martin Drive, Baltimore, MD 21218

G. FRITZ BENEDICT, BARBARA MCARTHUR, IVAN RAMIREZ, AND WILLIAM SPIESMAN

McDonald Observatory, University of Texas, Austin, TX 78712

AND

BURTON F. JONES

Lick Observatory, University of California, Santa Cruz CA 95064

(Accepted)

Draft version February 2, 2008

ABSTRACT

We present absolute trigonometric parallaxes and relative proper motions for three members of the Pleiades star cluster, obtained with *HST*'s Fine Guidance Sensor 1r, a white-light interferometer. We estimate spectral types and luminosity classes of the stars comprising the astrometric reference frame from $R \approx 2,000$ spectra, *VJHK* photometry, and reduced proper motions. From these we derive estimates of absolute parallaxes and introduce them into our model as observations with error. We constrain the three cluster members to have a 1σ dispersion in distance less than 6.4 pc, and find an average $\pi_{abs} = 7.43 \pm 0.17 \pm 0.20$ milliseconds of arc, where the second error is systematic due to member placement within the cluster. This parallax corresponds to a distance of 134.6 ± 3.1 pc or a distance modulus of $(m - M) = 5.65 \pm 0.05$ for these three Pleiads, presuming a central location. This result agrees with three other independent determinations of the Pleiades distance. Presuming that the cluster depth systematic error can be significantly reduced because of the random placement of these many members within the cluster, these four independent measures yield a best-estimate Pleiades distance of $\pi_{abs} = 7.49 \pm 0.07$ milliseconds of arc, corresponding to a distance of 133.5 ± 1.2 pc or a distance modulus of $(m - M) = 5.63 \pm 0.02$. This resolves the dispute between the main sequence fitting and the *Hipparcos* distance moduli in favor of main sequence fitting.

Subject headings: astrometry — interferometry — stars: distances — stars: clusters — distance scale

1. THE PROBLEM

Our knowledge of the life histories of stars relies on models whose fidelity is ultimately tested by appeal to real stars. The Sun provides the most basic calibration of these models, of course, because it is only for the Sun that an accurate age exists and for which the mass, temperature, composition, and structure are known with precision, accuracy, and completeness. Clusters of stars are also fundamental for constructing models because we can assume that all the cluster's members are of the same age and composition, even if other parameters are more loosely constrained.

Preeminent among clusters is the Pleiades, and much effort has gone into determining the absolute parallax of this cluster. ESA's *Hipparcos* mission brought the benefits of space observing to astrometry to produce precise positions, proper motions, and parallaxes for nearly all

stars brighter than $V \approx 9$. Before *Hipparcos*, the distance to the Pleiades was too large for ground-based parallaxes to yield a good distance, so the best estimates were derived by comparing the main sequence of the Pleiades to a main sequence constructed from nearby stars with large parallaxes. A small correction for evolution is necessary (the Pleiades is about 100 Myr old (Pinsonneault et al. 1998) while the nearby field stars are typically as old as the Sun), but the Pleiades appears to have essentially the same elemental abundances as the Sun (Boesgaard & Friel 1990), obviating a need for a metallicity correction such as is needed, for example, for the Hyades.

In addition to its primary program, *Hipparcos* included stars in several of the nearest open clusters in order to resolve the "Hyades distance problem" once and for all, and to similarly calibrate other clusters. The result obtained by *Hipparcos* for the Pleiades (van Leeuwen 1999) was a complete surprise, yielding a distance modulus of $(m - M) = 5.37 \pm 0.06$ magnitude, to be compared to a modulus of 5.60 ± 0.04 from main sequence fitting (Pinsonneault et al. 1998). Taken at face value, the *Hipparcos* result means that stars in the Pleiades are about 0.23 magnitude fainter than otherwise similar stars of the solar neighborhood. This large discrepancy has forced a careful reexamination of the assumptions and input parameters of the stellar models, as well as a thorough study of the *Hipparcos* data itself and potential errors

*BASED ON OBSERVATIONS MADE WITH THE NASA/ESA HUBBLE SPACE TELESCOPE, OBTAINED AT THE SPACE TELESCOPE SCIENCE INSTITUTE, WHICH IS OPERATED BY THE ASSOCIATION OF UNIVERSITIES FOR RESEARCH IN ASTRONOMY, INC., UNDER NASA CONTRACT NAS5-26555.

Electronic address: soderblom, nelan@stsci.edu

Electronic address: fritz, mca, ivan, spies@astro.as.utexas.edu

Electronic address: jones@ucolick.org

² Operated by the Association of Universities for Research in Astronomy, Inc. under contract to the National Aeronautics and Space Administration.

TABLE 1
PLEIADES LOG OF OBSERVATIONS AND MEMBER
PHOTOMETRY

Set	MJD	Roll ^a (degrees)	V ^b		
			3030	3063	3179
1	51770.65507	284.046	14.03	13.54	10.05
2	51783.6811	284.046	14.00	13.56	10.06
3	51957.17546	103.014	14.00	13.47	10.08
4	51968.37565	103.014	14.01	13.57	10.09
5	52128.77383	284.046	13.97	13.58	10.07
6	53053.24519	113.021	14.00	13.44	10.08
$\langle V \rangle$			14.00	13.53	10.07
σ_V			0.02	0.06	0.01

^aSpacecraft roll as defined in Chapter 2, FGS Instrument Handbook (Nelan & Makidon 2001)

^bAverage of 2 to 5 observations at each epoch. Internal errors are on order 0.005 magnitude per observation set.

in it. The controversy has not been fully resolved in that builders of star models find that the changes in physics or input parameters needed to account for the *Hipparcos* distance are too radical to be reasonable while the *Hipparcos* team has resolutely defended the *Hipparcos* result. With no clear reconciliation of these divergent views, we felt it worthwhile to reobserve some stars in the Pleiades in the traditional method of parallax astrometry—highly precise measurements of stellar positions relative to nearby reference stars—by taking advantage of the extraordinary precision achievable with Fine Guidance Sensor 1r on the *Hubble Space Telescope*.

This project began as an effort to resolve known Pleiades spectroscopic binaries into visual binaries so that we could both obtain an accurate distance and calibrate the Zero-Age Main Sequence with known masses. We did not succeed in resolving the spectroscopic binaries, nor would our measurement of the Pleiades parallax by itself resolve the problem raised by *Hipparcos*, but our measurement in concert with other recent independent measurements of the Pleiades distance clearly and unambiguously shows that the *Hipparcos* parallax is wrong and that traditional main sequence fitting results in reliable estimates. To avoid repetition, we will discuss the work to date in detail in our discussion.

2. OBSERVATIONS AND DATA REDUCTION

Six sets of astrometric data were acquired with *HST*, spanning 3.51 years, for a total of 135 measurements of the three Pleiads and nine reference stars. The three Pleiades targets were H II 3030, 3063, and 3179, identified hereafter by their Hertzsprung (1947) numbers. Table 1 lists the epochs of observation and measured FGS V-band photometry of the three Pleiads. Each data set required approximately 33 minutes of spacecraft time. The reductions and calibrations are detailed in Benedict et al. (2002a), Benedict et al. (2002b), and McArthur et al. (2001). At each epoch we measured both the reference stars and the target multiple times in order to correct for intra-orbit drift of the type seen in the cross filter calibration data shown in Figure 1 of Benedict et al. (2002a). Figure 1 shows the distribution of the reference stars (4–14) and the presumed




FIG. 1.— The Pleiades cluster, its members, and the astrometric reference stars observed.

Pleiads (3030, 3063, and 3179) on a second-generation *R*-band image, obtained from the Digital Sky Survey (<http://stdatu.stsci.edu/dss/>).

Bradley et al. (1991) and Nelan & Makidon (2001) provide an overview of *HST*'s Fine Guidance Sensors, and Benedict et al. (2002b) describe the fringe tracking (POS) mode astrometric capabilities of an FGS, along with data acquisition and reduction strategies also used in the present study. Times of observation use a modified Julian Date, $\text{MJD} = \text{JD} - 2444000.5$.

We obtained observations at each of the two maximum parallax factors. This leads to the two distinct spacecraft roll angles shown which result from the requirement to keep *HST*'s solar panels fully illuminated throughout the year. This roll constraint generally imposes alternate orientations at each time of maximum positive or negative parallax factor over a typical 2.5 year parallax campaign, allowing a clean separation of parallax and proper motion signatures. As noted, our original intent was to determine orbital parameters for some known spectroscopic binaries, but once resolution of the binary did not work out we changed this dynamical parallax experiment to a standard parallax program. The most recent data set extended our time span by 2.5 years, significantly improving the accuracy of our final parallaxes and the precision of our final proper motion values.

3. ABSOLUTE PARALLAXES FOR THE REFERENCE STARS

Because the parallax determined for the three Pleiades members is measured with respect to reference frame stars which have their own parallaxes, we must either apply a statistically-derived correction from relative to absolute parallax (van Altena, Lee, & Hoffleit (1995), hereafter YPC95), or estimate the absolute parallaxes of the reference frame stars. In principle, the colors, spectral type, and luminosity class of a star can be used to estimate the absolute magnitude, M_V , and V-band absorption, A_V . The absolute parallax is then simply,

$$\pi_{\text{abs}} = 10^{-(V-M_V+5-A_V)/5} \quad (1)$$




FIG. 2.— $(J - K)$ vs. $(V - K)$ color-color diagram. The dashed line is the locus of dwarf (luminosity class V) stars of various spectral types; the dot-dashed line is for giants (luminosity class III). The reddening vector indicates $A_V = 1.0$ for the plotted color systems.

The luminosity class is generally more difficult to estimate than the spectral type (temperature class), yet the derived absolute magnitudes are critically dependent on the assumed luminosity. As a consequence, we use as much additional information as possible in an attempt to confirm the luminosity classes. Specifically, we obtained 2MASS photometry and UCAC2 proper motions for a one-degree-square field containing Figure 1, and then iteratively employ the technique of reduced proper motion (Yong & Lambert (2003), Gould & Morgan (2003)) in an effort to discriminate between giants and dwarfs.

3.1. Reference Star Photometry

Our bandpasses for reference star photometry include: V (from FGS 1r), and JHK (from 2MASS¹). The 2MASS JHK values have been transformed to the Bessell & Brett (1988) system using the transformations provided in Carpenter (2001). Table 2 lists $VJHK$ photometry for the target and reference stars indicated in Figure 1.

3.2. Reference Star Spectroscopy

The spectra from which we estimated spectral type and luminosity class come from Lick Observatory². The resolution was approximately, with coverage from 3900 Å to 6700 Å. Classifications used a combination of template matching and line ratios. Spectral types for the stars are good to about 2 subclasses. Table 4 lists the spectral types and luminosity classes for our reference stars. The estimated classification uncertainties are used to generate the σ_{M_V} values in that table.

3.3. Interstellar Extinction

To determine interstellar extinction, we first plot these stars in a $(J - K)$ vs. $(V - K)$ diagram. A comparison

of the relationships between spectral type and intrinsic color against those we measured provides an estimate of reddening. Figure 2 shows this color-color diagram and a reddening vector for $A_V = 1.0$. Also plotted are mappings between spectral type and luminosity classes V and III from Bessell & Brett (1988) and Cox (2000) (hereafter AQ2000). Figure 2, along with the estimated spectral types, provides an indication of the reddening for each reference star.

Assuming an $R = 3.1$ Galactic reddening law (Savage & Mathis 1979), we derive A_V values by comparing the measured colors (Table 2) with intrinsic $(V - K)_0$ colors from Bessell & Brett (1988) and AQ2000. Specifically, we estimate A_V from $A_V/E(V - K) = 1.1$, derived from the Savage & Mathis (1979) reddening law. The resulting A_V values are collected in Table 3. Colors and spectral types of these reference stars are consistent with a field-wide average $\langle A_V \rangle = 0.17 \pm 0.06$, far less than the maximum reddening, $A_V < 0.72$, determined by Schlegel et al. (1998). For the stars classified as dwarfs, $\langle A_V \rangle = 0.14 \pm 0.03$, in good agreement with a recent determination of $A_V = 0.12$ for the Pleiades (Hainline et al. 2001). The more distant non-dwarfs have $\langle A_V \rangle = 0.23 \pm 0.08$.

The technique of reduced proper motions can confirm the reference stars' estimated luminosity classes, but the precision of existing proper motions for all the reference stars was so low that only suggestive discrimination between giants and dwarfs was possible. Typical uncertainties for H_K , a parameter equivalent to absolute magnitude, M_V , were about a magnitude. Nonetheless, a reduced proper motion diagram does suggest that ref-6, -9, and -10 are not dwarf stars, since they are considerably redder in $(J - K)$ than the other stars classified as dwarfs. Giants are typically redder in $(J - K)$ than dwarfs for a given spectral type (AQ2000). Our luminosity class uncertainty is reflected in their input spectrophotometric parallax errors (Table 4). We will revisit this additional test in Section 4.1, once we have solved for higher precision proper motions.

3.4. Adopted Reference Frame Absolute Parallaxes

We derive absolute parallaxes with M_V values from AQ2000 and the $\langle A_V \rangle$ derived from the photometry. Our adopted errors for $(m - M)_0$ are 0.7 mag for the dwarfs and from 0.7 to 2 mag for the non-dwarf reference stars. These are somewhat larger than we have used in the past (Benedict et al. (2002a), Benedict et al. (2002b), McArthur et al. (2002)), but justified given our far smaller set of spectrophotometric data. Our parallax values are listed in Table 4. Individually, no reference star absolute parallax is better determined than $\frac{\sigma_\pi}{\pi} = 32\%$. The average absolute parallax for the reference frame is $\langle \pi_{abs} \rangle = 1.3$ mas. As a check, we compare this to the correction to absolute parallax discussed and presented in YPC95 (Sec. 3.2, Fig. 2). Entering YPC95, Fig. 2, with the Pleiades Galactic latitude, $b = -23^\circ 0'$, and average magnitude for the reference frame, $\langle V_{ref} \rangle = 14.5$, we obtain a correction to absolute of 1.0 mas. We prefer to introduce into our reduction model our spectrophotometrically-estimated reference star parallaxes as observations with error. When such data are available, the use of spectrophotometric parallaxes offers a more direct (i.e., less Galaxy model-dependent) way of

¹ The Two Micron All Sky Survey is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology

² Lick Observatory is owned and operated by the University of California.

TABLE 2
FGS AND NEAR-IR PHOTOMETRY

ID	V	K	$(J - H)$	$(J - K)$	$(V - K)$
3179	10.07	8.68 ± 0.02	0.32 ± 0.02	0.37 ± 0.02	1.40 ± 0.10
3063	13.54	10.34 ± 0.02	0.67 ± 0.03	0.84 ± 0.03	3.20 ± 0.10
3030	14.00	10.63 ± 0.02	0.71 ± 0.03	0.85 ± 0.03	3.37 ± 0.10
ref-4	15.66	13.98 ± 0.05	0.43 ± 0.05	0.49 ± 0.06	1.70 ± 0.11
ref-6	14.56	12.03 ± 0.02	0.56 ± 0.02	0.68 ± 0.03	2.54 ± 0.10
ref-8	14.48	12.91 ± 0.03	0.29 ± 0.03	0.37 ± 0.04	1.57 ± 0.10
ref-9	13.60	10.64 ± 0.02	0.68 ± 0.02	0.79 ± 0.03	2.97 ± 0.10
ref-10	15.85	13.40 ± 0.04	0.55 ± 0.04	0.67 ± 0.05	2.45 ± 0.11
ref-11	14.63	12.75 ± 0.02	0.43 ± 0.03	0.49 ± 0.04	1.88 ± 0.10
ref-12	14.23	12.15 ± 0.03	0.48 ± 0.04	0.51 ± 0.04	2.10 ± 0.10
ref-13	13.15	10.57 ± 0.02	0.34 ± 0.03	0.42 ± 0.03	1.57 ± 0.10
ref-14	15.48	13.78 ± 0.03	0.42 ± 0.05	0.44 ± 0.05	1.70 ± 0.11

TABLE 3
REFERENCE STAR A_V FROM SPECTROPHOTOMETRY

ID	SpT	$(V - K)_0$	$(V - K)$	$E(V - K)$	A_V^a
ref-4	G5V	1.55	1.68	0.13	0.14
ref-6	K1IV	2.32	2.54	0.22	0.24
ref-8	G3V	1.45	1.57	0.12	0.13
ref-9	K2III	2.70	2.97	0.27	0.30
ref-10	K1IV	2.32	2.45	0.13	0.14
ref-11	G8V	1.80	1.88	0.08	0.09
ref-12	K0V	1.96	2.10	0.13	0.15
ref-13	G3V	1.45	1.57	0.12	0.13
ref-14	G5V	1.55	1.70	0.15	0.16

$$^a A_V = 1.1E(V - K)$$

TABLE 4
ASTROMETRIC REFERENCE STAR SPECTRAL CLASSIFICATIONS
AND SPECTROPHOTOMETRIC PARALLAXES

ID	Sp. T.	V	M_V	A_V	m-M	$\pi_{abs}(\text{mas})$
ref-4	G5V	15.68	5.1	0.14	10.6 ± 0.7	0.8 ± 0.3
ref-6	K1IV	14.5	3.4	0.23	11.2 ± 2	0.06 ± 0.6
ref-8	G3V	14.48	4.8	0.14	9.7 ± 0.7	1.2 ± 0.4
ref-9	K2III	13.61	0.5	0.23	13.1 ± 0.7	0.3 ± 0.1
ref-10	K1IV	15.85	3.4	0.23	12.5 ± 2	0.4 ± 0.3
ref-11	G8V	14.63	5.6	0.14	9.1 ± 0.7	1.7 ± 0.5
ref-12	K0V	14.24	5.9	0.14	8.3 ± 0.7	2.3 ± 0.7
ref-13	G3V	12.14	4.8	0.14	7.3 ± 0.7	3.66 ± 1.2
ref-14	G5V	15.48	5.1	0.14	10.4 ± 0.7	0.9 ± 0.3

determining the reference star absolute parallaxes.

4. THE ABSOLUTE PARALLAX OF THE PLEIADES

4.1. The Astrometric Model

Using the positions measured by FGS 1r, we determine the scale, rotation, and offset “plate constants” relative to an arbitrarily-adopted constraint epoch (the so-called “master plate”) for each observation set (the data acquired at each epoch). The MJD of each observation set is listed in Table 1, along with a measured magnitude transformed from the FGS instrumental system as per Benedict et al. (1998). Our Pleiades reference frame contains 9 stars. We employ the six-parameter model discussed in Benedict et al. (1999) for those observations.

In this case, we determined the plate parameters from target and reference star data. Additionally, we apply corrections for lateral color discussed in Benedict et al. (1999), using values specific to FGS 1r as determined from observations with that FGS.

As for all our previous astrometric analyses, we employ GaussFit (Jefferys et al. (1987)) to minimize χ^2 . The solved equations of condition for the Pleiades field are:

$$x' = x + lcx(B - V) \quad (2)$$

$$y' = y + lcy(B - V) \quad (3)$$

$$\xi = Ax' + By' + C - \mu_x \Delta t - P_\alpha \pi_x \quad (4)$$

$$\eta = -Bx' + Ay' + F - \mu_y \Delta t - P_\delta \pi_y \quad (5)$$

where x and y are the measured coordinates from *HST*; lcx and lcy are the lateral color corrections; and $(B - V)$ represents the $(B - V)$ color of each star, estimated from its spectral type, A_V , and $(J - K)$ color listed in Table 2. A and B are scale- and rotation plate constants, C and F are offsets; μ_x and μ_y are proper motions; Δt is the epoch difference from the mean epoch; P_α and P_δ are parallax factors; and π_x and π_y are the parallaxes in x and y . We obtain the parallax factors from a JPL Earth orbit predictor (Standish 1990), upgraded to version DE405. Orientation to the sky is obtained from ground-based astrometry (2MASS Catalog) with uncertainties in the field orientation of 0°05.

4.2. Modeling Constraints from Prior Knowledge

In addition to introducing our estimated reference star parallaxes as observations with error, we also introduce proper motion data from UCAC2 (Zacharias et al. (2003)) and Schilbach et al. (1995). Initial values are listed in Table 6. Typical input errors are 5-6 mas for each coordinate. The lateral color calibrations and the $(B - V)$ color indices are also treated as observations with error. As a final constraining observation, we solve for a line-of-sight dispersion in the parallaxes of the three Pleiades members with the ‘observation’ derived from the 1σ angular extent of the Pleiades (1° , from Adams et al. 2001) and an assumption of spherical symmetry. From this, we infer that the 1σ dispersion in distance in this group is $1^\circ/1 \text{ radian} = 1.7\%$. Hence, the 1σ dispersion in the parallax difference between Pleiades members is

$$\Delta\pi = 1.7\% \times \sqrt{2} \times 7.7 \text{ mas} = 0.20 \text{ mas}, \quad (6)$$




FIG. 3.— Reduced proper motion diagram for 5,542 stars in a one-degree field centered on the Pleiades. Star identifications are shown for our Pleiades targets (H II 3030, 3063, and 3179) and our astrometric reference stars. For a given spectral type, giants and sub-giants have more negative H_K values and are redder than dwarfs in $(J - K)$. Stars ref-6 and ref-10 are coincident. H_K values are derived from ‘Final’ proper motions in Table 6. The small cross at the lower left represents a typical $(J - K)$ error of 0.04 mag and H_K error of 0.17 mag. The horizontal dashed line is a giant-dwarf demarcation derived from a statistical analysis of the Tycho input catalog (Ciardi 2004, private communication).

where we have here temporarily adopted a parallax of the Pleiades, $\langle \pi \rangle = 7.7$ mas. The parallax dispersion among targets 3030, 3179, and 3063 becomes an observation with associated error fed to our model, an observation used to estimate the parallax dispersion among the three stars, while solving for their parallaxes. Loosening the cluster 1σ dispersion to 2° (i.e., $\Delta\pi = 0.38$ mas) had no effect on the final weighted average parallax. Again, note that $\Delta\pi = 0.2$ mas is not an error associated with the *distance* to the Pleiades. It serves to constrain the dispersion in distances measured for Pleiades members.

Proper motion values obtained from our modeling of *HST* data are listed in Table 6 as ‘Final’. We now employ the technique of reduced proper motions to provide a confirmation of the reference star estimated luminosity class listed in Table 4. We obtain proper motion and J , K photometry from UCAC2 and 2MASS for a one-degree-square field centered on the Pleiades. Figure 3 shows $H_K = K + 5 \log(\mu)$ versus $(J - K)$ color index for 5,542 stars. If all stars had the same transverse velocities, Figure 3 would be equivalent to an H-R diagram. Target Pleiads and reference stars are plotted as ID numbers from Table 6. Errors in H_K are now ~ 0.3 mag. Reference stars 6, 9, and 10 are clearly separated from the others, supporting their classification as non-dwarfs. Ref-6 and ref-10 remain below ref-9, confirming their sub-giant nature.

4.3. Assessing Reference Frame Residuals

Our initial modeling attempts indicated that three of the original twelve reference stars exhibited significantly larger residuals than average. These reference stars were




FIG. 4.— Histograms of x and y residuals obtained from modeling the Pleiades members and astrometric reference stars with equations 4 and 5. Distributions are fit with gaussians whose 1σ dispersions are noted in the plots.

TABLE 5
PLEIADES AND REFERENCE STAR POSITIONS

ID	V	ξ^a	η^a
3179	10.08	$+163.1991 \pm 0.0002$	-13.1667 ± 0.0002
3063	13.47	-198.8822 ± 0.0003	$+59.7779 \pm 0.0003$
3030	14.00	-268.2952 ± 0.0003	$+39.2897 \pm 0.0003$
ref-4	15.68	$+213.0875 \pm 0.0012$	$+51.9487 \pm 0.0010$
ref-6	14.57	$+99.5992 \pm 0.0003$	$+51.2020 \pm 0.0006$
ref-8 ^b	14.47	0.0000 ± 0.0005	0.0000 ± 0.0006
ref-9	13.61	-8.4395 ± 0.0003	-42.3189 ± 0.0003
ref-10	15.85	-60.3818 ± 0.0009	-26.3444 ± 0.0008
ref-11	14.63	-292.6740 ± 0.0003	$+28.8663 \pm 0.0004$
ref-12	14.24	-254.6792 ± 0.0004	$+109.6633 \pm 0.0005$
ref-13	12.14	-156.6322 ± 0.0003	$+97.8026 \pm 0.0003$
ref-14	15.48	-338.2213 ± 0.0006	-19.8165 ± 0.0007

^a ξ and η are relative positions in arcseconds

^bRA = 03 51 45.050, Dec = +23 53 43.43, J2000

near the top- and bottom edges of the FGS1r field of regard. The Optical Field Angle Distortion calibration (McArthur et al. (2002)) reduces as-built *HST* telescope and FGS1r distortions with amplitude $\sim 1''$ to below 2 mas over much of the FGS1r field of regard. However, because the fidelity of correction drops precipitously near the edge of the field of regard, we removed these three




FIG. 5.— Absolute parallax determinations for the Pleiades. We compare astrometric parallax results (filled diamonds) from *HST* (“*HST*”), *Hipparcos* (“*Hip*”), and recent determinations from Allegheny Observatory (“*AO*”), Gatewood, de Jonge, & Han (2000). Pan, Shao & Kulkarni (2004) (“*Pan*”) have derived a dynamical parallax from long-baseline interferometry and radial velocity measurements of the binary star Atlas. Munari et al. (2004) have performed a similar dynamical determination on another Pleiades binary. MS denotes a parallax derived from main-sequence fitting (Pinsonneault et al. 1998). The horizontal dashed line is the weighted average of the *HST*, Pan, Munari and AO measures.

stars from the solution. From histograms of the remaining reference star astrometric residuals (Figure 4) we conclude that we have obtained satisfactory correction. The resulting reference frame ‘catalog’ in ξ and η standard coordinates (Table 5) was determined with $\langle\sigma_\xi\rangle = 0.5$ and $\langle\sigma_\eta\rangle = 0.5$ mas.

To determine if there might be unmodeled—but possibly correctable—systematic effects at the 1 mas level, we plotted the Pleiades reference frame x and y residuals against a number of spacecraft, instrumental, and astronomical parameters. These included x and y position within the FGS “pickle”; radial distance from the pickle center; reference star V magnitude and $(B-V)$ color; and epoch of observation. We saw no obvious trends, other than an expected increase in positional uncertainty with reference star magnitude.

4.4. The Absolute Parallax of The Pleiades

Note that we do not measure the parallax of these Pleiads relative to a reference frame with unknown parallax and then apply a correction to absolute parallax, assuming some model of the Galaxy. In a quasi-Bayesian approach, the reference star spectrophotometric absolute parallaxes, UCAC2 and Schilbach et al. (1995) proper motions, and an estimated cluster depth were input as observations with associated errors, not as hard-wired quantities known to infinite precision. Parallaxes and relative proper motion results from *HST* are collected in Tables 6 and 7. We obtain for the Pleiades members an average absolute parallax $\pi_{abs} = 7.43 \pm 0.17$ mas (Table 8). Because we employ a cluster depth constraint, the three Pleiades member parallaxes are not independent measurements. Hence, we cannot use the standard deviation of the mean to reduce our final error by $\sqrt{2}$. Along with our result, other recent Pleiades parallaxes are listed in Table 9 and compared in Figure 5. The most discrepant of these is clearly and only the *Hippar-*




FIG. 6.— Bottom: *HST* absolute parallax determinations compared with *Hipparcos* for all targets listed in Table 10. Top: The *Hipparcos* residuals to the dotted error-weighted impartial regression line that excludes the Pleiades. The error bars on the residuals are *Hipparcos* Catalog 1σ errors.

cos result.

Our absolute parallax for the Pleiades contains one last systematic uncertainty: where in the cluster do our three Pleiades members lie? In Section 4.2 we estimated a ‘depth’ in parallax of ~ 0.20 mas. Our final parallax result should be stated $\pi_{abs} = 7.43 \pm 0.17 \pm 0.20$ mas with the error having both a random and systematic component. We point out that each of the astrometric results in Figure 5 suffers from the same systematic error. In the next Section we reduce that error by averaging those results. Inspecting Tables 6 and 7 we note that ref-14, identified as Cl* Melotte 22 CALAR 7, is in fact not a Pleiad, disagreeing in parallax and proper motion with the first three stars in these Tables, all identified members.

5. DISCUSSION AND SUMMARY

5.1. *HST* Parallax Accuracy

Our parallax precision, an indication of our internal, random error, is often less than 0.3 mas. To assess our accuracy, or external error, we must compare our parallaxes with results from independent measurements. Following Gatewood, Kiewiet de Jonge, & Persinger (1998) and extending the analysis presented in Benedict et al. (2002b) with the addition of a recent parallax for Gl 876 (Benedict et al. 2002c), we plot eight parallaxes obtained by the *HST* Astrometry Science Team with FGS 3 and, now FGS 1r, against those obtained by *Hipparcos*. Data for these objects are collected in Table 10 and shown in Figure 6. The dashed line is a weighted regression that takes into account errors in both input data sets and excludes the Pleiades. Figure 6 indicates no statistically significant scale difference compared to *Hipparcos*. However, for this fit, which excludes the Pleiades, we obtain a reduced $\chi^2 = 0.265$. Including the Pleiades, we ob-

TABLE 6
PLEIADES AND REFERENCE STAR PROPER MOTIONS

ID	V	Input (UCAC2)		Final (<i>HST</i>)	
		μ_x^a	μ_y^a	μ_x^a	μ_y^a
3179	10.08	$+0.0192 \pm 0.0006$	-0.0465 ± 0.0006	$+0.0192 \pm 0.0003$	-0.0465 ± 0.0002
3063 ^b	13.47	$+0.0164 \pm 0.0011$	-0.0418 ± 0.0011	$+0.0168 \pm 0.0003$	-0.0421 ± 0.0005
3030 ^b	14.00	$+0.0154 \pm 0.0004$	-0.0408 ± 0.0004	$+0.0155 \pm 0.0004$	-0.0403 ± 0.0002
ref-4	15.68	$+0.0035 \pm 0.0056$	-0.0034 ± 0.0056	$+0.0024 \pm 0.0026$	-0.0019 ± 0.0024
ref-6	14.57	$+0.0054 \pm 0.0056$	-0.0094 ± 0.0056	$+0.0044 \pm 0.0010$	-0.0049 ± 0.0010
ref-8	14.47	$+0.0052 \pm 0.0056$	-0.0119 ± 0.0056	$+0.0030 \pm 0.0005$	-0.0016 ± 0.0005
ref-9	13.61	$+0.0117 \pm 0.0056$	$+0.0033 \pm 0.0056$	-0.0014 ± 0.0015	$+0.0031 \pm 0.0019$
ref-10	15.85	-0.0016 ± 0.0069	-0.0092 ± 0.0069	-0.0016 ± 0.0022	-0.0091 ± 0.0026
ref-11	14.63	$+0.0030 \pm 0.0056$	-0.0131 ± 0.0056	-0.0043 ± 0.0007	-0.0039 ± 0.0006
ref-12	14.24	$+0.0058 \pm 0.0056$	-0.0096 ± 0.0056	-0.0008 ± 0.0010	$+0.0024 \pm 0.0012$
ref-13	12.14	-0.0074 ± 0.0019	-0.0129 ± 0.0019	-0.0093 ± 0.0010	-0.0083 ± 0.0011
ref-14	15.48	0.0000 ± 0.0058	-0.0053 ± 0.0058	-0.0056 ± 0.0024	-0.0051 ± 0.0032

^a μ_x and μ_y are relative motions in arcsec yr⁻¹

^b μ_x and μ_y from Schilbach et al. (1995).

TABLE 7
PLEIADES AND REFERENCE STAR
PARALLAXES AND TRANSVERSE VELOCITIES

ID	μ^a mas yr ⁻¹	π_{abs}^b mas	V_t^c km s ⁻¹
3179	50.36 ± 0.40	7.45 ± 0.16	32
3063	45.30 ± 0.53	7.43 ± 0.16	29
3030	43.20 ± 0.48	7.41 ± 0.18	28
4	3.07 ± 3.54	0.82 ± 0.09	18
6	6.63 ± 1.39	0.84 ± 0.25	38
8	3.40 ± 0.76	1.21 ± 0.13	13
9	3.42 ± 2.43	0.26 ± 0.03	61
10	9.28 ± 3.39	0.36 ± 0.11	122
11	5.79 ± 0.98	1.66 ± 0.16	17
12	2.50 ± 1.52	2.25 ± 0.23	5
13	12.50 ± 1.46	1.64 ± 0.32	36
14	7.58 ± 4.04	0.92 ± 0.10	39

^a $\mu = (\mu_x^2 + \mu_y^2)^{1/2}$ from ‘Final’ in Table 6

^b Final π_{abs} from modeling *HST* data with equations 2–5, employing the constraints summarized in Section 4.2

^c $V_t = 4.74 \times \mu / \pi_{abs}$

TABLE 8
PLEIADES PARALLAX AND PROPER MOTION

Parameter	Value
<i>HST</i> study duration	3.51 y
number of observation sets	6
reference star $\langle V \rangle$	14.63
reference star $\langle (B - V) \rangle$	0.9 ^a
<i>HST</i> Absolute Parallax ^b	7.43 ± 0.17 mas
<i>HST</i> Relative Proper Motion ^c in pos. angle	46.3 ± 3.7 mas y ⁻¹ $158^\circ \pm 1^\circ$

^a Estimated from *VJHK* photometry and spectral types, with $A_V = 0.14$ for dwarfs and $A_V = 0.23$ for giants.

^b Average of 3030, 3063, and 3179 from Table 7

^c Average of 3030, 3063, and 3179 from Table 6, ‘Final.’ Proper motion error is the standard deviation of the individual measures.

tain a significantly poorer fit with reduced $\chi^2 = 0.551$, again, suggesting a problem with the *Hipparcos* Pleiades parallax.

Our result, in and of itself, does not lead to the conclusion that the *Hipparcos* parallax for the Pleiades is wrong, but that conclusion cannot be avoided once all the results are examined together. Especially important for making this case are the two recent determinations of visual binary orbits for Pleiades members. Pan, Shao & Kulkarni (2004) used the Palomar Testbed Interferometer to determine very precise relative positions of the two stars comprising Atlas, one of the Seven Sisters. Without having a radial velocity orbit they could not determine all the parameters, but a solution is possible by assuming masses for the stars, and the masses enter in the cube root. By doing this they concluded that the distance to the Pleiades cannot be less than 127 pc and that the most likely distance lies between 133 and 137 pc. Munari et al. (2004) analyzed light- and radial velocity curves for HD 23642, an eclipsing binary in the Pleiades, and determined a distance of 132 ± 2 pc. (This would decrease to 130.6 ± 3.7 if the assumed reddening were increased to as much as $E(B - V) = 0.035$ magnitude).

5.2. The Distance to the Pleiades

There now exist three completely independent determinations of the Pleiades distance that use completely independent techniques and data, and they all yield the same answer to within their errors. Our traditional parallax determination leads to $d = 134.6 \pm 3.1$ pc, a visual binary orbit leads to $d = 135 \pm 2$, and an eclipsing binary orbit results in $d = 132 \pm 2$. For comparison, recent estimates from main sequence fitting include 132 ± 4 (Stello & Nissen 2001), and 132 ± 2 (Pinsonneault et al. 1998), and Gatewood, de Jonge, & Han (2000) has determined 131 ± 7 at Allegheny Observatory. Narayanan & Gould (1999) derived 131 ± 24 pc from the gradient in the radial velocities of Pleiades members in the direction of the cluster’s proper motion.

Clearly the *Hipparcos* result, 118 ± 4 , is discrepant. This can be seen graphically in Figure 6, and a summary of these distance determinations is given in

TABLE 9
PREVIOUS AND PRESENT PLEIADES PARALLAXES

Method	Abbr.	π_{abs}	d (pc)	$(m - M)$	Reference
<i>HST</i> /FGS parallax	<i>HST</i>	7.43 ± 0.17	134.6 ± 3.1	5.65 ± 0.05	this paper
Hipparcos all-sky	HIP	8.45 ± 0.25	118.3 ± 3.5	5.37 ± 0.06	van Leeuwen (1999)
Allegheny Obs.	AO	7.64 ± 0.43	130.9 ± 7.4	5.59 ± 0.11	Gatewood, de Jonge, & Han (2000)
interferometric orbit	Pan	7.41 ± 0.11	135.0 ± 2.0	5.65 ± 0.03	Pan, Shao & Kulkarni (2004)
dynamical parallax	Mun	7.58 ± 0.11	131.9 ± 3.0	5.60 ± 0.05	Munari et al. (2004)
main sequence fitting	MS	7.58 ± 0.14	131.9 ± 2.4	5.60 ± 0.04	Pinsonneault et al. (1998)

TABLE 10
HST AND *Hipparcos* ABSOLUTE PARALLAXES

Object	π_{HST} (mas)	π_{HIP} (mas)	<i>HST</i> Reference
Proxima Cen	769.7 ± 0.3	772.33 ± 2.42	Benedict et al. (1999)
Barnard's Star	545.5 ± 0.3	549.3 ± 1.58	Benedict et al. (1999)
Gliese 876	214.6 ± 0.2	212.7 ± 2.1	Benedict et al. (2002c)
Feige 24	14.6 ± 0.4	13.44 ± 3.62	Benedict et al. (2000)
Wolf 1062	98.0 ± 0.4	98.56 ± 2.66	Benedict et al. (2001)
Pleiades	7.43 ± 0.17	8.45 ± 0.25	this paper
RR Lyrae	3.60 ± 0.20	4.38 ± 0.59	Benedict et al. (2002a)
δ Cephei	3.66 ± 0.15	3.32 ± 0.58	Benedict et al. (2002b)
HD 213307	3.65 ± 0.15	3.43 ± 0.64	Benedict et al. (2002b)

Table 9. Understanding this discrepancy is crucial. Astrometry is arguably the one branch of astronomy where accurate and precise knowledge of uncertainties cannot be overlooked. The *Hipparcos* team has been well aware of this and has put considerable effort into examining potential sources of systematic error. Their most recent papers (van Leeuwen & Fantino (2003a), Dalla Torre & van Leeuwen (2003), van Leeuwen & Penston (2003), Fantino & van Leeuwen (2003), van Leeuwen & Fantino (2003b)) show, for instance, that noise in the along-scan attitude dominates for $H_p < 4.5$ (where H_p is the apparent magnitude as directly measured by *Hipparcos*) and that this may be especially important for the Pleiades, inter alia (van Leeuwen & Fantino 2003b). This possibility was examined by Makarov (2002), who reanalyzed *Hipparcos* data to derive $d = 129 \pm 3$, a value that is substantially less discrepant than that reported by van Leeuwen (1999) and Robichon et al. (1999).

The answer certainly does not lie in an unusual shape or physical properties for the Pleiades. Stello & Nissen (2001) suggested that the *Hipparcos* distance could be reconciled with traditional measures if the bright stars—the Seven Sisters—that dominate the *Hipparcos* result happen to lie at the near end of an elongated cluster. This is disproved by the fact that Pan, Shao & Kulkarni (2004) find Atlas itself to lie at the traditional distance. Grenon (2001) suggested that the luminosities of Pleiades stars could be accounted for by a low cluster metallicity of -0.112 ± 0.025 , determined from Geneva photometry. The exact metallicity of the Pleiades remains uncertain, but it is unlikely to be as low as that since analyses from high-resolution spectra yield values that are essentially solar (e.g., Boesgaard & Friel (1990) get $[\text{Fe}/\text{H}] = -0.034 \pm 0.024$). Hainline et al. (2001) have likewise refuted the Grenon (2001) metallicity on several grounds.

To summarize, *HST* astrometry yields an absolute trigonometric parallax for three members of the Pleiades, $\pi_{abs} = 7.43 \pm 0.17$ mas with a 0.20 mas systematic error due to cluster depth. A weighted average with previous ground-based astrometric determinations (*HST*, AO, Pan and Munari, Table 9) provides $\pi_{abs} = 7.49 \pm 0.07$ mas. This average result should reduce the contribution of the cluster depth systematic error, presuming that the stars measured by these techniques are randomly distributed within the cluster. With $\frac{\sigma_\pi}{\pi} \sim 1\%$, any Lutz-Kelker-Hanson bias correction (Lutz & Kelker (1973), Hanson (1979)) to an absolute magnitude would be less than 0.01 magnitude (e.g., Benedict et al. (2002b)). This net parallax of 7.49 ± 0.07 mas corresponds to $d = 133.5 \pm 1.2$ pc, or $(m - M) = 5.63 \pm 0.02$ magnitude. This is likely to be the best available distance for the Pleiades until observations of substantially better precision can be made with a mission such as SIM or GAIA.

We thank an anonymous referee for comments which materially improved the clarity of our presentation. Support for this work was provided by NASA through grant GO-08335 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. These results are based partially on observations obtained with the Lick Observatory 3 m telescope, which is owned and operated by the University of California. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by NASA and the NSF. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France; the NASA/IPAC Extragalactic Database (NED) which is operated by JPL,

California Institute of Technology, under contract with the NASA; and NASA's Astrophysics Data System Ab-

stract Service.

REFERENCES

- Adams, J. D., Stauffer, J. R., Monet, D. G., Skrutskie, M. F., & Beichman, C. A. 2001, *AJ*, 121, 2053
- Allen, C. W. 1973, London: University of London, Athlone Press, 3rd ed.
- van Altena, W. F., Lee, J. T., & Hoffleit, E. D. 1995, *Yale Parallax Catalog* (4th ed. ; New Haven, CT: Yale Univ. Obs.) (YPC95)
- Benedict, G. F., McArthur, B., Nelan, E., Story, D., Whipple, A. L., Shelus, P. J., Jefferys, W. H., Hemenway, P. D., Franz, O. G., Wasserman, L. H., Duncombe, R. L., van Altena, W., & Fredrick, L. W. 1998, *AJ*, 116, 429
- Benedict, G. F., McArthur, B., Chappell, D. W., Nelan, E., Jefferys, W. H., van Altena, W., Lee, J., Cornell, D., Shelus, P. J., Hemenway, P. D., Franz, O. G., Wasserman, L. H., Duncombe, R. L., Story, D., Whipple, A., & Fredrick, L. W. 1999, *AJ*, 118, 1086
- Benedict, G. F., McArthur, B. E., Franz, O. G., Wasserman, L. H., Nelan, E., Lee, J., Fredrick, L. W., Jefferys, W. H., van Altena, W., Robinson, E. L., Spiesman, W. J., Shelus, P. J., Hemenway, P. D., Duncombe, R. L., Story, D., Whipple, A. L., & Bradley, A. 2000, *AJ*, 119, 2382
- Benedict, G. F., McArthur, B. E., Franz, O. G., Wasserman, L. H., Henry, T. J., Takato, T., Strateva, I. V., Crawford, J. L., Ianna, P. A., McCarthy, D. W., Nelan, E., Jefferys, W. H., van Altena, W., Shelus, P. J., Hemenway, P. D., Duncombe, R. L., Story, D., Whipple, A. L., Bradley, A. J., & Fredrick, L. W. 2001, *AJ*, 121, 1607
- Benedict, G. F., McArthur, B. E., Fredrick, L. W., Harrison, T. E., Lee, J., Slesnick, C. L., Rhee, J., Patterson, R. J., Nelan, E., Jefferys, W. H., van Altena, W., Shelus, P. J., Franz, O. G., Wasserman, L. H., Hemenway, P. D., Duncombe, R. L., Story, D., Whipple, A. L., & Bradley, A. J. 2002a, *AJ*, 123, 473
- Benedict, G. F., McArthur, B. E., Fredrick, L. W., Harrison, T. E., Slesnick, C. L., Rhee, J., Patterson, R. J., Skrutskie, M. F., Franz, O. G., Wasserman, L. H., Jefferys, W. H., Nelan, E., van Altena, W., Shelus, P. J., Hemenway, P. D., Duncombe, R. L., Story, D., Whipple, A. L., & Bradley, A. J. 2002b, *AJ*, 124, 1695
- Benedict, G. F., McArthur, B. E., Forveille, T., Delfosse, X., Nelan, E., Butler, R. P., Spiesman, W., Marcy, G., Goldman, B., Perrier, C., Jefferys, W. H., & Mayor, M. 2002c, *ApJ*, 581, L115
- Bessell, M. S. 1979, *PASP*, 91, 5
- Bessell, M. S. & Brett, J. M. 1988, *PASP*, 100, 1134
- Boesgaard, A. M., & Friel, E. D. 1990, *ApJ*, 351, 467
- Bradley, A., Abramowicz-Reed, L., Story, D., Benedict, G. & Jefferys, W. 1991, *PASP*, 103, 317
- Carpenter, J. M. 2001, *AJ*, 121, 2851
- Cox, A. N. 2000, *Allen's Astrophysical Quantities*, 4th ed. Publisher: New York: AIP Press, Springer, 2000. Edited by Arthur N. Cox. (AQ2000)
- Dalla Torre, A., & van Leeuwen, F. 2003, *Sp. Sci. Rev.*, 108, 451
- Fantino, E., & van Leeuwen, F. 2003, *Sp. Sci. Rev.*, 108, 499
- Gatewood, G., Kiewiet de Jonge, J., & Persinger, T. 1998, *AJ*, 116, 1501
- Gatewood, G., de Jonge, J. K., & Han, I. 2000, *ApJ*, 533, 938
- Gould, A. & Morgan, C. W. 2003, *ApJ*, 585, 1056
- Grenon, M. 2001, in *11th Cool Stars, Stellar Systems, and the Sun*, eds. R. J. García López, R. Rebolo, and M. R. Zapatero Osorio, A.S.P. Conf. Ser., v. 223, 359
- Hainline, L. J., Deliyannis, C. P., & Margheim, S. 2001, *BAAS*, 33, 1387
- Hanson, R. B. 1979, *MNRAS*, 186, 87
- Harrison, E. 1947, *Ann. Leiden Obs.*, 19, Part 1A
- Jefferys, W., Fitzpatrick, J., and McArthur, B. 1987, *Celest. Mech.* 41, 39.
- van Leeuwen, F. 1999, *A&A*, 341, L71
- van Leeuwen, F. 2002, in *Highlights of Astron.*, ed. H. Rickman, 12, 657
- van Leeuwen, F., & Fantino, E. 2003, *Sp. Sci. Rev.*, 108, 447
- van Leeuwen, F., & Fantino, E. 2003, *Sp. Sci. Rev.*, 108, 537
- van Leeuwen, F., & Penston, M. 2003, *Sp. Sci. Rev.*, 108, 471
- Lutz, T. E. & Kelker, D. H. 1973, *PASP*, 85, 573
- Makarov, V. V. 2002, *AJ*, 124, 3299
- McArthur, B., Benedict, G. F., Jefferys, W. H., & Nelan, E. 2002, *The 2002 HST Calibration Workshop : Hubble after the Installation of the ACS and the NICMOS Cooling System*, Proceedings of a Workshop held at the Space Telescope Science Institute, Baltimore, Maryland, October 17 and 18, 2002. Edited by Santiago Arribas, Anton Koekemoer, and Brad Whitmore. Baltimore, MD: Space Telescope Science Institute, 2002, p.373
- McArthur, B. E. et al. 2001, *ApJ*, 560, 907
- Mermilliod, J.-C., Rosvick, J. M., Duquennoy, A., & Mayor, M. 1992, *A&A*, 265, 513
- Munari, U., Dallaporta, S., Siviero, A., Soubiran, C., Fiorucci, M., and Girard, P. 2004, *A&A*, 418, L31
- Narayanan, V. K., & Gould, A. 1999, *ApJ*, 523, 328
- Nelan, E., & Makidon, R., in *HST FGS Data Handbook*, version 4.0, ed. B. Mobasher, Baltimore, STScI
- Nelan, E., et al., "Fine Guidance Sensor Instrument Handbook," version 12.0, Baltimore, STScI
- Nelan, E., & Makidon, R. 2001, *Fine Guidance Sensor Instrument Handbook* (version 10; Baltimore: STScI)
- Pan, X., Shao, M., & Kulkarni, S., 2004, *Nature*, 427, 326
- Percival, S. M., Salaris, M., & Kilkenny, D. 2003, *A&A*, 400, 541
- Pinsonneault, M. H., Stauffer, J., Soderblom, D. R., King, J. R., & Hanson, R. B. 1998, *ApJ*, 504, 170
- Robichon, N., Arenou, F., Mermilliod, J.-C., & Turon, C. 1999, *A&A*, 345, 471
- Savage, B. D. & Mathis, J. S. 1979, *ARA&A*, 17, 73
- Schilbach, E., Robichon, N., Souchay, J., & Guibert, J. 1995, *A&A*, 299, 696
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Soderblom, D. R., King, J. R., Hanson, R. B., Jones, B. F., Fischer, D., Stauffer, J. R., & Pinsonneault, M. H. 1998, *ApJ*, 504, 192
- Soderblom, D. R., Stauffer, J. R., Hudon, J. D., & Jones, B. F. 1998, *ApJS*, 85, 315
- Standish, E. M., Jr. 1990, *A&A*, 233, 252
- Stello, D. & Nissen, P. E. 2001, *A&A*, 374, 105
- Yong, D. & Lambert, D. L. 2003, *PASP*, 115, 796
- Zacharias, N., Urban, S. E., Zacharias, M. I., Wycoff, G. L., Hall, D. M., Germain, M. E., Holdenried, E. R., & Winter, L. 2003, *VizieR Online Data Catalog*, 1289